

## Angiosperm symbioses with non-mycorrhizal fungal partners enhance N acquisition from ancient organic matter in a warming maritime Antarctic

Hill, Paul; Broughton, Richard; Bougoure, Jeremy; Havelange, William; Newsham, Kevin K.; Gran, Helen; Murphy, Daniel V.; Clode, Peta; Ramayah, Soshila; Marsden, Karina; Quilliam, Richard; Roberts, Paula; Brown, Caley; Read, David J.; DeLuca, Thomas; Bardgett, Richard D.; Hopkins, David W.; Jones, Davey L.

### Ecology Letters

DOI:  
[10.1111/ele.13399](https://doi.org/10.1111/ele.13399)

Published: 01/12/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

#### *Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Hill, P., Broughton, R., Bougoure, J., Havelange, W., Newsham, K. K., Gran, H., Murphy, D. V., Clode, P., Ramayah, S., Marsden, K., Quilliam, R., Roberts, P., Brown, C., Read, D. J., DeLuca, T., Bardgett, R. D., Hopkins, D. W., & Jones, D. L. (2019). Angiosperm symbioses with non-mycorrhizal fungal partners enhance N acquisition from ancient organic matter in a warming maritime Antarctic. *Ecology Letters*, 22(12), 2111-2119. <https://doi.org/10.1111/ele.13399>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Angiosperm symbioses with non-mycorrhizal fungal partners enhance N acquisition from ancient organic matter in a warming maritime Antarctic**

Paul W Hill<sup>1</sup>, Richard Broughton<sup>2,3</sup>, Jeremy Bougoure<sup>4,5</sup>, William Havelange<sup>1</sup>, Kevin K Newsham<sup>2</sup>, Helen Grant<sup>6</sup>, Daniel V Murphy<sup>4</sup>, Peta Clode<sup>5,7</sup>, Soshila Ramayah<sup>1</sup>, Karina A Marsden<sup>1,8</sup>, Richard S. Quilliam<sup>1,9</sup>, Paula Roberts<sup>1</sup>, Caley Brown<sup>1</sup>, David J Read<sup>10</sup>, Thomas H Deluca<sup>1,11</sup>, Richard D Bardgett<sup>12</sup>, David W Hopkins<sup>13</sup>, Davey L Jones<sup>1,4</sup>

1 School of Natural Sciences, Bangor University, Bangor LL57 2UW, UK.

2 British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, UK.

3 Institute of Aquaculture, University of Stirling, Stirling, FK9 4LA, UK

4 SoilsWest, UWA School of Agriculture and Environment, Faculty of Science, University of Western Australia, WA 6009, Australia.

5 Centre for Microscopy, Characterisation and Analysis, University of Western Australia, WA 6009, Australia.

6 Life Sciences Mass Spectrometry Facility, Lancaster Environment Centre, Lancaster LA1 4AP UK.

7 UWA School of Biological Sciences, University of Western Australia, WA 6009, Australia.

8 Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, Victoria 3010, Australia.

9 Biological and Environmental Sciences, Faculty of Natural Sciences, University of Stirling, Stirling, FK9 4LA, UK.

10 Animal and Plant Sciences, University of Sheffield, Western bank, Sheffield, S10 2TN, UK

11 WA Franke College of Forestry and Conservation, University of Montana, Missoula, MT 59812 USA.

27 12 School of Earth and Environmental Sciences, University of Manchester, Manchester, M13  
28 9PL, UK.

29 13 SRUC - Scotland's Rural College, West Mains Road, Edinburgh, EH9 3JG, UK.

30

31

32 Running title: DSEs enhance Antarctic plant N acquisition

33 Keywords: Soil; Nitrogen cycle; Carbon cycle; Climate change; Enantiomers; Polar; Dark  
34 septate endophytes

35 Type or article: Letter

36 Words in Abstract: 150

37 Words in Main text: 3676

38 Number of references: 40

39 Number of figures: 5

40 Corresponding author:

41 Paul Hill, Environment Centre Wales, School of Natural Sciences, Bangor University, Deiniol  
42 Road, Bangor LL57 2UW, UK.

43 Email: p.w.hill@bangor.ac.uk

44 Tel: +44 1248 2632

45 Author Contributions:

46 PH, DJ, KKN, RDB, DH, PR, TD and RQ conceived the investigation; PH carried out  
47 fieldwork; RB carried out amino acid analysis; JB, DM and PC carried out nanoSIMS work;  
48 PH, WH, CB, SR and KM carried out laboratory experiments and analysis; HG carried out  
49 IRMS analysis; PH wrote the manuscript first draft; all authors contributed to the final version.

50 Data accessibility:

51 Data will be archived at Figshare DOI: 10.6084/m9.figshare.9791693 following acceptance.

52

## Abstract

In contrast to the situation in plants inhabiting most of the world's ecosystems, mycorrhizal fungi are usually absent from roots of the only two native vascular plant species of maritime Antarctica, *Deschampsia antarctica* and *Colobanthus quitensis*. Instead, a range of ascomycete fungi, termed dark septate endophytes (DSEs), frequently colonise the roots of these plant species. We demonstrate that colonisation of Antarctic vascular plants by DSEs facilitates not only the acquisition of organic nitrogen as early protein breakdown products, but also as non-proteinaceous D-amino acids and their short peptides, accumulated in slowly-decomposing organic matter, such as moss peat. Our findings suggest that, in a warming maritime Antarctic, this symbiosis has a key role in accelerating the replacement of formerly dominant moss communities by vascular plants, and in increasing the rate at which ancient carbon stores laid down as moss peat over centuries or millennia are returned to the atmosphere as CO<sub>2</sub>.

## Introduction

Fungal root symbionts have been crucial to the success of plants in terrestrial ecosystems, with a relationship dating back to the colonisation of land (Strullu-Derrien et al. 2018). Mutualistic relationships with mycorrhizal fungi remain key to the acquisition of limiting nutrients, such as nitrogen (N) and phosphorus (P), in the majority of terrestrial plants (Smith & Read 2008). However, in marked contrast to their presence in most ecosystems, mycorrhizas are typically absent from the roots of vascular plants in maritime Antarctica (Upson et al. 2008; Newsham et al. 2009). In this region, the roots of the two native angiosperms, *Deschampsia antarctica* Desv. (a grass) and *Colobanthus quitensis* (Kunth) Bartl. (a cushion-forming plant, Fig. 1) are instead colonised by a range of ascomycete fungi, collectively termed dark septate endophytes (DSEs) (Fig. 1, Upson et al. 2008; Newsham et al. 2009), which may have a role in the acquisition of organic N from soils (Upson et al. 2009; Newsham 2011).

In areas of the maritime Antarctic not under permanent ice, moss cover can be extensive (Fig. S1 in Supporting Information) and dominates primary productivity. It is estimated to account for 45 km<sup>2</sup> of Antarctic Peninsula land area and is particularly prevalent on islands such as the South Orkney and South Shetland Islands (Fretwell et al. 2011; Royles & Griffiths 2015). Due to the constraints imposed on decomposition by low temperatures, moss growth leads to the accumulation of large amounts of soil organic matter, including substantial stores of protein (Royles & Griffiths 2015). Vascular plants, and particularly pioneer individuals and populations, are commonly found amongst mosses, exploiting stored proteinaceous N to facilitate establishment (Fig. 1, Hill et al. 2011a).

In a survey of roots of *D. antarctica* and *C. quitensis* on Signy Island (60° 43' S, 45° 36' W) in the South Orkney Islands, maritime Antarctica, we found the most consistent and extensive occurrence of DSE hyphae and characteristic microsclerotia (Fig. 1) was in the roots of plants growing amongst banks formed by the moss *Chorisodontium aciphyllum* (Hook. f. & Wilson) Broth. (Fig. 1). Banks formed by this moss frequently exceed 1 m in depth and may be up to 3 m deep, storing organic matter that has remained undecomposed over millennia (Royles et al. 2012; Royles & Griffiths 2015, Fig. S2 in Supporting Information). This organic matter has become increasingly bioavailable as mean air temperatures have risen in the maritime Antarctic, leading to progressive thawing of the moss banks (Royles et al. 2012; Abrams et al. 2013; Royles & Griffiths 2015; Amesbury et al. 2017).

In most cases (e.g., amongst the moss *Sanionia uncinata* (Hedw.) Loeske, Figs. 1 and S1), *D. antarctica* appears to root no deeper than c. 10 cm, with its roots usually extending to a depth of 5 cm or less (Fig. S3 in Supporting Information), corresponding to the depth of accumulated organic matter. However, in *C. aciphyllum* banks, the grass was observed rooting down to > 25 cm, where organic matter may have been stored for > 500 years (Royles et al., 2012). We hypothesised that the penetration of roots colonised by DSEs deep into moss banks allows *D.*

*antarctica* to exploit ancient nutrients that up until recent decades were unavailable because the moss banks have been frozen.

Due to slow N mineralisation, it is likely that early breakdown products of accumulated proteins (L-amino acids and their short peptides) make a substantial contribution to plant N nutrition in polar soils (Chapin et al. 1993; Hill et al. 2011a). However, peptides containing D-glutamic acid and especially D-alanine are common constituents of bacterial peptidoglycan and various D-amino acids occur in bacteria, archaea, fungi, plants and animals (Yoshimura & Esaki 2003; Friedman 2010; Vranova et al. 2012). D-amino acids are also known to accumulate from proteinaceous L-amino acids during long periods of storage, due to abiotic racemisation, which may take place at a rates of about 0.3% of L-amino acids per decade (Wichern et al. 2004). Consequently, D-amino acids accumulate in soils where decomposition is slow e.g., in deserts or in peat, such as that formed by moss banks (Kunnas & Jauhiainen 1993; Wichern et al. 2004).

It is clear from previous investigations that both plants and soil microbes are able to take up and metabolise some D-amino acids such as D-alanine (Hill et al. 2011b,c; Hill et al. 2012; Vranova et al. 2012). However, in contrast to short L-peptides, which appear to be widely metabolised, until now, evidence suggested that short D-peptides could be metabolised by soil microbes but not by plants (Hill et al. 2011b,c; Hill et al. 2012; Vranova et al. 2012). Whether the ability to metabolise D-peptides is present in plants inhabiting soils where D-enantiomers are a more available source of N is unknown. We measured uptake of a range of N forms under field conditions in the Antarctic and found that both native vascular plants could acquire N from D-alanine and its dipeptide - as well as from longer peptides of the L-enantiomer than previously recognised. Further, we found that colonisation with DSEs facilitated plant acquisition of N from both L- and D-enantiomers of alanine and their peptides.

## **Materials and Methods**

### **Assessment of fungal endophyte colonisation**

Roots of *D. antarctica* and *C. quitensis* were collected from locations around Signy Island (Gourlay Peninsula; Polynesia Point; Factory Cove; Berntsen Point; Lower slopes of Factory Bluffs; Starfish Cove; North Point; Moss Braes; Deschampsia Point; Foca Cove; Fig. S4 in Supporting Information). Roots were washed in water and examined for the presence of DSE hyphae and microsclerotia by light microscopy after staining (Newsham & Bridge 2010). The same analyses confirmed the absence of arbuscular mycorrhizal structures from roots (Upson et al. 2008).

### **Soil solution collection**

Rhizon soil solution samplers (5 cm long; Rhizosphere Research Products, Wageningen, Netherlands) were inserted into soil under mosses (mostly *S. uncinata* and *C. aciphyllum*) or vascular plants (*D. antarctica* with some *C. quitensis*). Soil solution was sampled over a depth of c. 2–6 cm at approximately fortnightly intervals for about 12 weeks during austral summer. Large soluble proteins and peptides were then removed by passing solutions through a 1 kDa ultrafiltration membrane (Millipore, Billerica, MA, USA).

### **Analysis of amino acid enantiomers**

Filtered soil solution samples taken over the season from each site were pooled, divided in two and concentrated by freeze drying. One portion was hydrolysed for 16 h in 6 M HCl under N<sub>2</sub> and freeze-dried again. The dry soil solution residues were re-suspended in 500 µl of 0.01 M HCl with 1.875 pmol µl<sup>-1</sup> of L-homoarginine as the internal standard. Amino acid enantiomers were quantified by HPLC (Broughton et al. 2015).

### **Substrate uptake in intact plant-soil system**

Monoliths (c. 20 × 20 cm) of *D. antarctica* or *C. quitensis* growing in native soil were collected from the Moss Braes region of Signy Island and stored outside for about 24 h prior to

experiments. About 1–2 h prior to experiments, 15 mm diameter, 40 mm deep plugs were taken from the monoliths. Solutions (2.5 ml) of 98 at%  $^{15}\text{N}$  (inorganic) or dual  $^{15}\text{N}$ ,  $^{13}\text{C}$  (organic) 1 mM L-alanine, D-alanine, L-dialanine, D-dialanine, L-trialanine, L-tetraalanine, L-pentaalanine,  $\text{NH}_4\text{Cl}$  or  $\text{KNO}_3$  (L-enantiomers, and inorganic from CK-Gas Products, Hook, UK; D-enantiomers from Sigma-Aldrich, Gillingham, UK) were injected into plugs ( $n=4$  and  $n=3$  for *D. antarctica* or *C. quitensis*, respectively). After 1 h in daylight at *c.* 2 °C, shoot material was removed, dried (80 °C) and ground before analysis in a Eurovector Isoprime IRMS (Eurovector SpA, Milan, Italy).

#### **Sterile culture of *D. antarctica* and inoculation of roots with DSEs**

Sterile individuals of *D. antarctica* (we were not able to generate a sterile culture of *C. quitensis*) were prepared according to a protocol modified from Cuba et al. (2005). Plants were removed from soil and washed in tap water. Roots and shoots were trimmed and the remaining tissue was shaken in  $\text{NaHClO}_3$  (*c.* 14% free Cl) with 1 drop of Tween 20 for 25 min, followed by 80% ethanol for 5 min. After thorough washing in sterile tap water, remaining leaf and root was trimmed from crown tissue, which was then placed on the surface of sterile agar containing 2.1 g l<sup>-1</sup> Murashige & Skoog basal medium, 1 mmol l<sup>-1</sup> glucose and 47 μmol l<sup>-1</sup>  $\text{NaSiO}_3$  in Phytatrays (Sigma-Aldrich, Gillingham, UK). Amphotericin B solution (5 ml of 2.5 mg l<sup>-1</sup>) was then added to the surface of agar around the crown tissue. Plants were grown at 10°C with a 16 h photoperiod at *c.* 500 μmol photons m<sup>-2</sup> s<sup>-1</sup>. Tillers were separated periodically and replanted in agar as above (except for amphotericin B, which was not used after the first culture). Any Phytatrays showing signs of microbial contamination were discarded. Examination of roots of sterilised plants by light microscopy and TEM did not reveal the presence of any microbes.

Sterile plants for use in experiments were transplanted into Phytatrays containing sterile perlite with *c.* 100 ml of 2.1 g l<sup>-1</sup> Murashige & Skoog basal medium, 1 mmol l<sup>-1</sup> glucose and 47 μmol



$1^{-1}$  NaSiO<sub>3</sub> with and without inoculation with a DSE (*Tapesia* sp.; Helotiales; GenBank accession #FN178471) which was isolated from roots of *D. antarctica* growing on Coronation Island, around 7 km from where experimental plants and soils were collected. At least three weeks was allowed for the DSE to colonise roots before plants were used in experiments. Plants were then removed from the inoculated perlite and grown in uninoculated perlite, as used for the controls.

#### **Substrate uptake from sterile solution**

Sterile or DSE-inoculated *D. antarctica* plants were removed from perlite and roots gently washed in sterile 0.1 mM KCl, followed by de-ionised water. Roots of intact plants ( $n=4$ ) were then placed in sterile vials containing 2 ml of 100  $\mu$ M, 98 at% <sup>15</sup>N (inorganic) or dual <sup>15</sup>N, <sup>13</sup>C (organic) L-alanine, D-alanine, L-dialanine, D-dialanine, L-trialanine, D-trialanine, L-tetraalanine, L-pentaalanine, NH<sub>4</sub>Cl or KNO<sub>3</sub>. After 1 h, plants were removed from solutions, washed in de-ionised water followed by 100 mM CaCl<sub>2</sub>. Roots and shoots were separated and analysed by IRMS, as above.

#### **Plant metabolism of substrates**

To determine whether substrates could be metabolised, sterile or DSE-inoculated roots of intact *D. antarctica* plants ( $n=3$ ) were submerged in 2 ml of 10  $\mu$ M, c. 7.5 kBq ml<sup>-1</sup> 1-<sup>14</sup>C L-alanine, D-alanine, L-dialanine, D-dialanine, L-trialanine, D-trialanine, L-tetraalanine or L-pentaalanine (American Radiolabeled Chemicals, St Louis, MO, USA). Vials and plants were sealed in 50 ml clear polypropylene containers. Air was drawn through containers at 300 ml min<sup>-1</sup> and bubbled through 15 ml Oxysolve C-400 Scintillant (Zinsser Analytic, Frankfurt, Germany) to capture respired <sup>14</sup>CO<sub>2</sub>. Carbon dioxide traps were changed after 10, 20, 40, 60 and 80 min and captured <sup>14</sup>CO<sub>2</sub> measured by scintillation counting in a Wallac 1404 scintillation counter (Perkin-Elmer Life Sciences, Waltham, MA, USA).

202 After 80 min, plants were removed from solutions, washed as above and dried. Dry roots and  
203 shoots were combusted in a Harvey OX400 Biological Oxidiser (Harvey Instruments Corp.,  
204 Hillsdale, NJ, USA). Liberated  $^{14}\text{CO}_2$  was captured in Oxysolve C-400 and  $^{14}\text{C}$  activity  
205 measured by liquid scintillation counting as above.

#### 206 **Uptake kinetics**

207 Sterile or DSE-inoculated roots of intact *D. antarctica* plants ( $n=3$ ) were submerged in labelled  
208 ( $^{14}\text{C}$  or  $^{15}\text{N}$  for organic and inorganic substrates, respectively) substrate solutions as above. In  
209 this case, exposure to solutions was for 15 min and substrate concentrations were 1, 5, 10, 50,  
210 100, 250, 500, 750  $\mu\text{M}$  and 1, 2.5, 5, 7.5 and 10 mM. Plants were analysed for  $^{14}\text{C}$  or  $^{15}\text{N}$  as  
211 above. Respired  $^{14}\text{CO}_2$  was captured in Oxysolve C-400 and measured as above. Michaelis-  
212 Menten constants were calculated from hyperbolic fits to uptake data (Sigmaplot v13, Systat,  
213 Hounslow, UK).

#### 214 **NanoSIMS analysis**

215 Sterile or DSE colonised *D. antarctica* ( $n=3$ ) roots were submerged in 3 mM solution of either  
216  $^{13}\text{C}^{15}\text{N}$  D-trialanine or  $^{13}\text{C}^{15}\text{N}$  -L-trialanine. Plants were incubated for 5 mins then removed  
217 from isotope enriched solution, washed quickly in MQ water, then high pressure frozen (HPF;  
218 1 mm segments) in hexadecene cryoprotectant (EM PACT2, Leica Microsystems, Wetzlar,  
219 Germany). HPF samples were cryosubstituted (EM AFS2, Leica Microsystems, Wetzlar,  
220 Germany) using the method described in Bougoure et al. (2014). Briefly, samples were  
221 immersed in prechilled ( $-130\text{ }^{\circ}\text{C}$ ) acrolein:diethyl ether over molecular sieve and brought to  
222 room temperature over 3 weeks before being infiltrated and embedded in epoxy resin. Sections  
223 250 nm thick were cut dry (i.e. not floated onto water for collection), mounted on Si wafers,  
224 and Au coated (10 nm) for nanoSIMS analysis. Regions of interest were identified and imaged  
225 at 120 kV in a transmission electron microscope (TEM; JEOL 2100) fitted with a digital camera  
226 (Gatan, ORIUS1000; Gatan Inc., Pleasanton, CA, USA). Sections were also collected on glass

slides, stained with toluidine blue and examined by optical microscopy to guide locations of nanoSIMS analyses.

*In situ* isotopic mapping was done using a NanoSIMS 50 (Cameca, Gennevilliers, France), with a 16 keV Cs<sup>+</sup> primary ion beam. Analyses were performed in multi-collection mode simultaneously detecting negative secondary ions <sup>12</sup>C<sub>2</sub>, <sup>12</sup>C<sup>13</sup>C, <sup>12</sup>C<sup>14</sup>N, and <sup>12</sup>C<sup>15</sup>N. The mass spectrometer was tuned to high mass resolution of c. 10000 (CAMECA definition) to separate <sup>12</sup>C<sup>15</sup>N from <sup>13</sup>C<sup>14</sup>N using an entrance slit of 30 μm, an aperture slit of 200 μm, and a 10% reduction in the signal at the energy slit. For secondary ion imaging, the primary current was set to c. 2 pA using a 350-μm primary aperture, giving a spot size of c. 100 nm. Analyses were done in chain mode so individual 30 × 30 μm analyses (256 pixel resolution) could be montaged to generate a dataset across entire root sections. All areas were implanted to the same ion dose (6 × 10<sup>16</sup> ions cm<sup>-2</sup>) prior to each acquisition.

Images were processed using the OpenMIMS data analysis software (National Resource for Imaging Mass Spectrometry <http://nrims.harvard.edu>) for the freeware package ImageJ (National Institutes of Health, Bethesda, MD, USA). Images were corrected for detector dead time (44 ns) on individual pixels and montages were produced using NRRD mosaics script (<http://nrims.harvard.edu>).

#### **Statistical analyses**

Data were analysed by *t*-test, one-way ANOVA with Tukey HSD post-hoc test or repeated measures ANOVA (SPSS v22; IBM, New York, USA) after testing for normality and homogeneity of variance with Shapiro-Wilk and Levene's tests, respectively. Data not conforming were transformed prior to analysis. Where a suitable transformation could not be identified, Games-Howell test was used. Statistical differences were accepted at  $P \leq 0.05$  unless otherwise stated.

## Results

### Amino acid concentrations in soil solution

The presence of vascular plants was associated with increases ( $P \leq 0.05$ ) in soil solution concentrations of 16 out of 18 measured free amino acids (L-enantiomers and glycine) by as much as ten-fold compared to sites where mosses grew alone (Fig. 2). The concentrations of non-protein D-amino acids were more variable, but there was more than three times as much free D-alanine, D-glutamate, D-histidine and D-threonine ( $P \leq 0.05$ ) in soil with vascular plants compared to moss-only soil (the concentrations of three other D-amino acids were greater with statistical significance at  $P < 0.1$ ). Soluble, peptide-bound amino acids tended to be present in soil solution at concentrations approximately ten-times greater than free amino acids (statistically different at  $P \leq 0.05$  for 20 and 21 amino acid enantiomers under vascular plants and mosses, respectively). The concentrations of almost half of the bound L-amino acids and D-alanine and D-histidine were greater ( $P \leq 0.05$ ) when vascular plants were present, relative to mosses alone.

### Uptake of amino acids and peptides under field conditions

Tests of uptake of a range of N forms under field conditions in the Antarctic showed that both native vascular plant species could acquire  $^{15}\text{N}$  from D-alanine and its dipeptide - as well as from peptides of the L-enantiomer up to five amino acids in length (Fig. 3). Rates of uptake appeared similar between the two species. Recovery of amino acid and peptide  $^{13}\text{C}$  suggested some intact uptake of molecules, although lack of data for root material and losses of  $^{13}\text{C}$  in respiration prevented quantification (Fig. S5 in Supporting Information). Although DSEs were present in the roots of plants used in these experiments, whether the fungal endophytes influenced nutrient acquisition could not be established.

## **Uptake, partitioning and metabolism of amino acids and peptides by plants with sterile roots or colonised with DSEs**

Although there were minor differences between isotopic tracers, with the exception of nitrate, DSE colonisation increased the uptake of all forms of N supplied to roots, with strong positive effects of the endophyte on the uptake of L-tri-, L-tetra- and L-pentaalanine ( $P < 0.05$ ; Fig. 4). Nitrate was also the only tested form of N where Michaelis-Menten constants for N uptake showed no indication of an effect of DSE colonisation (Table S1 in Supporting Information). Surprisingly, the DSE appeared to promote N translocation such that colonised plants had a lower ratio of root  $^{15}\text{N}$  to shoot  $^{15}\text{N}$  than uninoculated control plants ( $P < 0.001$ ; Fig. S6 in Supporting Information). Further, in contrast to limited data for other plants, loss of  $^{14}\text{CO}_2$  in respiration demonstrated that *D. antarctica* could metabolise all forms of organic N supplied, including D-peptides (Fig. S7 in Supporting Information; Hill et al. 2011c). However, actual rates of C loss in respiration are probably somewhat overestimated due to the  $^{14}\text{C}$  label being located only on the carboxyl group (Dippold & Kuzyakov 2013; Hill & Jones 2019). Nanoscale Secondary Ion Mass Spectrometry (nanoSIMS) showed transfer of L-peptide  $^{15}\text{N}$  into the intercellular space between the root cortical cells of *D. antarctica* by DSE hyphae (Fig. 5; Fig. S8 in Supporting Information). Additionally, individual root cells of plants supplied with D- or L-trialanine were more enriched with  $^{15}\text{N}$  when colonised with the DSE than in sterile controls, strongly suggesting that enhanced isotope recovery in bulk root analyses was not merely separate uptake by roots and fungus.

## **Discussion**

It appears that the presence of vascular plants in the organic soils of the maritime Antarctic gives rise to a marked increase in availability of both L- and D-enantiomers of amino acids as N sources. This suggests a stimulation of the rate of breakdown of stored moss peat in the

301 presence of roots, probably resulting from rhizosphere priming (Gavazov et al. 2018). Of free  
302 (and peptide-bound) D-amino acids, D-alanine was amongst the most available, maintaining  
303 concentrations around 10% of those of L-alanine, despite microbial consumption at rates  
304 similar to those of L-amino acids, indicating a significant production flux in these soils (Hill et  
305 al. 2011b). Whether this D-alanine originates primarily from peptidoglycan, abiotic  
306 racemisation of L-alanine in stored proteins, or another process is currently unknown.  
307 Similarly, although we can attribute occurrence of other D-amino acids to racemisation, it is  
308 not clear whether this is the only or even the principal source (Vranova et al. 2012). However,  
309 irrespective of the exact origin, the actual increase in availability of amino acid-N driven by  
310 vascular plants is likely to be greater than the increase in measured soil solution concentrations,  
311 due to a probable higher consumption flux from both microbes and plant roots in soils under  
312 vascular plants than under mosses (Hill et al. 2011a,b).

313 DSEs are widespread in plant roots in a range of ecosystems (Jumpponen 2001; Newsham et  
314 al. 2008), but there has been limited identification of their roles in plant nutrient acquisition to  
315 date, with some appearing to have negative effects on plant hosts (Jumpponen 2001; Upson et  
316 al. 2009; Newsham 2011; Vergara et al. 2018). Consequently, it remains unknown whether  
317 symbioses with DSEs are widespread facilitators of nutrient acquisition. It is clear from the  
318 findings here that the colonisation of roots by DSEs has a marked effect on the ability of  
319 Antarctic angiosperms to exploit amino acid N. The nanoSIMS images demonstrate direct  
320 hyphal transfer of peptide N to the root, and the surprising effect of DSE colonisation on  
321 translocation of N suggests an additional physiological effect on the host plant (direct hyphal  
322 transfer to shoots is unlikely due to confinement of this group of fungi to roots; Rodriguez et  
323 al. 2009). Colonisation appears to aid acquisition of some forms of N, such as peptides of D-  
324 amino acids and an L-pentapeptide, which have not previously been recognised as viable  
325 sources of N for plants. This may be due to the probable higher availability of both L- and D-

enantiomers in ecosystems where large quantities of proteinaceous material accumulate and  
 turn over slowly (Chapin et al. 1993; Kunnas & Jauhiainen 1993; Wichern et al. 2004). The  
 occurrence of close relatives of the DSE used here in the Arctic may support this view  
 (Genbank accessions MF920427 and KF617231; Krishnan et al. 2018; Taylor et al. 2014).  
 However, as both D- and L-peptides do exist in other ecosystems and investigation into plant  
 use of D-peptide N has been limited, it may be that the use of these N forms by both plants and  
 DSEs is more widespread than is currently recognised (Friedman 2010; Hill et al. 2011c;  
 Vranova et al. 2012). Some mosses are also colonised by endophytic fungi, but there is no  
 evidence for a role of these endophytes in nutrient acquisition (Davey & Currah 2006).  
 As greenhouse gas emissions to the atmosphere continue, near-surface air temperatures in the  
 maritime Antarctic are projected to warm by 2–4 °C by 2100 (Bracegirdle et al. 2008). Our  
 measurements suggest that vascular plants could increase rates of organic matter breakdown  
 under Antarctic mosses by up to an order of magnitude. Rising air temperatures are known to  
 synergistically increase rhizosphere priming, with increases in temperature sensitivity of,  
 perhaps, 25-50% in the presence of living roots (Boone et al. 1998; Zhu & Cheng 2011; Hill et  
 al. 2015). Hence, it appears that priming of ancient organic matter stored in moss banks arising  
 from plant growth and warming may interact to further increase nutrient availability, enhancing  
 the proliferation of angiosperms and returning more C to the atmosphere in a complex positive  
 feedback (Convey & Smith 2006; Day et al. 2008; Cannone et al. 2016; Gavazov et al. 2018;  
 Newsham et al. 2018). Thus, it seems probable that the stocks of moss-derived organic matter  
 accumulated over millennia will disappear at increasingly rapid rates as temperatures rise and  
 the ecology of the maritime Antarctic changes.

## References

350 Abram, N.J. et al. (2013). Acceleration of snow melt in an Antarctic Peninsula ice core during  
 351 the twentieth century. *Nat. Geosci.*, 6, 404–411.

352 Amesbury, M.J. et al. (2017). Widespread biological response to rapid warming on the  
 353 Antarctic Peninsula. *Curr. Biol.*, 27, 1616-1622.

354 Boone, R.D., Nadelhoffer, K.J., Canary, J.D. & Kaye, J.P. (1998). Roots exert a strong  
 355 influence on the temperature sensitivity of soil respiration. *Nature*, 396, 570–572.

356 Bougoure, J. et al. (2014). High resolution secondary ion mass spectrometry analysis of carbon  
 357 dynamics in mycorrhizas formed by an obligately myco-heterotrophic orchid. *Plant Cell*  
 358 *Environ.*, 37, 1223-1230.

359 Bracegirdle, T.J., Connolley, W.M. & Turner, J. (2008). Antarctic climate change over the  
 360 twenty first century. *J. Geophys. Res.*, 113, D03103

361 Broughton, R.C.I., Newsham, K.K., Hill, P.W., Stott, A. & Jones, D.L. (2015). Comparison of  
 362 the incorporation of D- and L-enantiomeric forms of alanine and its peptides into PLFAs by  
 363 different components of an Antarctic soil microbial community. *Soil Biol. Biochem.*, 88, 83-  
 364 89.

365 Cannone, N., Guglielmin, M., Convey, P., Worland, M.R. & Favero Longo, S.E. (2016).  
 366 Vascular plant changes in extreme environments: effects of multiple drivers. *Clim. Change*,  
 367 134, 651-665.

368 Chapin, F.S., Moilanen, L. & Kielland, K. (1993). Preferential use of organic nitrogen for  
 369 growth by a non-mycorrhizal arctic sedge. *Nature*, 361, 150-153.

370 Convey, P. & Smith, R.I.L. (2006). Responses of terrestrial Antarctic ecosystems to climate  
 371 change. *Plant. Ecol.*, 182, 1-10.

372 Cuba, M., Gutiérrez-Moraga, A., Butendieck, B. & Gidekel, M. (2005). Micropropagation of  
 373 *Deschampsia antarctica* – a frost-resistant Antarctic plant. *Antarct. Sci.*, 17, 69-70.



374 Day, T.A., Ruhland, C.T. & Xiong, F.S. (2008). Warming increases aboveground plant  
 375 biomass and C stocks in vascular-plant-dominated Antarctic tundra. *Glob. Change Biol.*, 14,  
 376 1827-1843.

377 Dippold, M.A. & Kuzyakov, Y. (2013). Biogeochemical transformations of amino acids in soil  
 378 assessed by position-specific labelling. *Plant Soil*, 373, 385-401.

379 Friedman, M. (2010) Origin, microbiology, nutrition and pharmacology of D-amino acids.  
 380 *Chem. Biodivers.* 7, 1491-1530.

381 Gavazov, K. et al. (2018). Vascular plant-mediated controls on atmospheric carbon  
 382 assimilation and peat carbon decomposition under climate change. *Glob. Change Biol.*, 24,  
 383 3911-3921.

384 Hill, P.W. et al. (2015). Living roots magnify the response of soil organic carbon  
 385 decomposition to temperature in temperate grassland. *Glob. Change Biol.*, 21, 1368-1375.

386 Hill, P.W. et al. (2011a). Vascular plant success in a warming Antarctic may be due to efficient  
 387 nitrogen acquisition. *Nat. Clim. Change*, 1, 50-53.

388 Hill, P.W., Farrell, M. & Jones, D.L. (2012). Bigger may be better in soil N cycling: does rapid  
 389 acquisition of small L-peptides by soil microbes dominate fluxes of protein-derived N in soil?  
 390 *Soil Biol. Biochem.*, 48, 106-112.

391 Hill, P.W. et al. (2011b). Soil- and enantiomer-specific metabolism of amino acids and their  
 392 peptides by Antarctic soil microorganisms. *Soil Biol. Biochem.*, 43, 2410-2416.

393 Hill, P.W. et al. (2011c). Acquisition and assimilation of nitrogen as peptide-bound and D-  
 394 enantiomers of amino acids by wheat. *PLoS ONE*, 6, e19220.

395 Hill, P.W. & Jones, D.L. (2019). Plant-microbe competition: does injection of isotopes of C  
 396 and N into the rhizosphere effectively characterise plant use of soil N? *New Phytol.* 221, 796-  
 397 806.

398 Jumpponen, A. (2001). Dark septate endophytes – are they mycorrhizal? *Mycorrhiza*, 11, 207-  
399 211.

400 Krishnan, A., Convey, P., Gonzalez, M., Smykla, J. & Alias, S.A. (2018). Effects of  
401 temperature on extracellular hydrolase enzymes from soil microfungi. *Polar Biol.*, 41, 537-  
402 551.

403 Kunnas, A.V. & Jauhiainen, T.-P. (2011). Separation and identification of free amino acid  
404 enantiomers in peat by capillary gas chromatography. *J. Chromatogr.* **628**, 269-273 (1993).

405 Newsham, K.K. A meta-analysis of plant responses to dark septate root endophytes. *New*  
406 *Phytol.*, 190, 783–793.

407 Newsham, K. K. & Bridge, P.D. (2010). Sebacinaleae are associates of the leafy liverwort  
408 *Lophozia excisa* in the southern maritime Antarctic. *Mycorrhiza*, 20, 307–313.

409 Newsham, K.K., Garnett, M.H., Robinson, C.H. & Cox, F. (2018). Discrete taxa of  
410 saprotrophic fungi respire different ages of carbon from Antarctic soils. *Sci. Rep.*, 8, 7866.

411 Newsham, K.K., Upson, R. & Read, D.J. (2009). Mycorrhizas and dark septate root endophytes  
412 in polar regions. *Fungal Ecol.*, 2, 10-20.

413 Rodriguez, R.J., White, J.F., Arnold, A.E. & Redman, R.S. (2009). Fungal endophytes:  
414 diversity and functional roles. *New Phytol.* 182, 314-330.

415 Royles, J. & Griffiths, H. (2015). Climate change impacts in polar regions: lessons from  
416 Antarctic moss bank archives. *Glob. Change Biol.*, 21, 1041-1057.

417 Royles, J. et al. (2012). Carbon isotope evidence of recent climate-related enhancement of CO<sub>2</sub>  
418 assimilation and peat accumulation rates in Antarctica. *Glob. Change Biol.*, 18, 3112-3124.

419 Smith, S.E. & Read, D.J. (2008). *Mycorrhizal Symbiosis*. Academic press.

420 Strullu-Derrien, C., Selosse, M.-A., Kenrick, P. & Martin, F.M. (2018). The origin and  
421 evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytol.*,  
422 220, 1012-1030.

Taylor, L.D., Hollingsworth, T.N., McFarland, J.W., Lennon, N.J., Nusbaum, C. & Ruess, R.W. (2014). A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning. *Ecol. Monogr.*, 84, 3-20.

Upton, R., Newsham, K.K. & Read, D.J. (2008). Root-fungal associations of *Colobanthus quitensis* and *Deschampsia antarctica* in the Maritime and Subantarctic. *Arct. Antarct. Alp. Res.*, 40, 592-599.

Upton, R., Read, D.J. & Newsham, K.K. (2009). Nitrogen form influences the response of *Deschampsia antarctica* to dark septate root endophytes. *Mycorrhiza*, 20, 1-11.

Vergara, C. et al. (2018). Dark septate endophytic fungi increase green manure-<sup>15</sup>N recovery efficiency, N contents and micronutrients in rice grains. *Front. Plant Sci.*, 9, 613.

Vranova, V. et al. (2012). The significance of D-amino acids in soil, fate and utilization by microbes and plants: review and identification of knowledge gaps. *Plant Soil*, 354, 21-39.

Wichern, F. et al. (2004). Changes in amino acid enantiomers and microbial performance in soils from a subtropical mountain oasis in Oman abandoned for different periods. *Biol. Fert. Soils*, 39, 398-406.

Yoshimura, T. & Esaki, N. (2003). Amino acid racemases; functions and mechanisms. *J. Biosci. Bioeng.* 96, 103-109.

Zhu, B. & Cheng, W. (2011). Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition. *Glob. Change Biol.*, 17, 2172-2183.

## Supporting Information

Additional Supporting Information may be downloaded via the online version of this article at Wiley Online Library ([www.ecologyletters.com](http://www.ecologyletters.com)).

## Acknowledgements

449 We thank Stacey Adlard and Bruce Maltman for assistance with fieldwork, Liam Smallwood,  
450 Peter Talboys and Sarah Chesworth for laboratory assistance, Rebecca Upson and British  
451 Antarctic Survey staff who are too numerous to mention by name. Special thanks to Jonathan  
452 and Linda Roberts. The authors acknowledge the facilities of Microscopy Australia at the  
453 Centre for Microscopy, Characterisation & Analysis, The University of Western Australia, a  
454 facility funded by the University, State and Commonwealth Governments. This work was  
455 funded by UK Natural Environment Research Council grant NE/I012303/1.

456

## Figures


**Figure 1** Antarctic vascular plants exploiting areas previously colonised by mosses on Signy Island and DSEs in roots of *Deschampsia antarctica*. a. *D. antarctica* growing in a bank of *Chorisodontium aciphyllum*. b. *D. antarctica* growing through mixed *Sanionia uncinata* and *Polytrichum juniperinum*. c. *D. antarctica* growing amongst *Andreaea* sp. d. *Colobanthus quitensis* growing through *C. aciphyllum*. e. *C. quitensis* growing through *S. uncinata*. f. *D. antarctica* and *C. quitensis* growing with *S. uncinata*. g. DSE hyphae in *D. antarctica* root. h. DSE microsclerotium (arrowed) in *D. antarctica* root (scale bars on panels g and h are 20  $\mu$ m).

**Figure 2** Concentrations of D- and L-enantiomers of amino acids in soil solutions at Signy Island under mosses alone or where vascular plants are present. a. free amino acids. b. amino acids bound in soluble peptides. Values are means  $\pm$  SEM;  $n=23$  and  $n=16$  for free and bound amino acids, respectively, under vascular plants;  $n=26$  and  $n=21$  for free and bound amino acids, respectively, under mosses only. Asterisks indicate differences between soil where vascular plants are present or where mosses are present alone ( $P \leq 0.05$ ).


**Figure 3** Rates of uptake of inorganic N and D- and L-enantiomers of alanine and short peptides thereof into shoots of *D. antarctica* and *C. quitensis* following injection of  $^{15}\text{N}$ - and  $^{13}\text{C}$ -labelled substrates into soil. Values are mean  $\pm$  SEM;  $n=3$  or 4.

**Figure 4** Rates of uptake by *D. antarctica* of N supplied in different forms. N uptake calculated from recovery of  $^{14}\text{C}$  (a) and  $^{15}\text{N}$  (b). Data are mean  $\pm$  SEM;  $n=3$  and  $n=4$  for  $^{14}\text{C}$  and  $^{15}\text{N}$ , respectively. Calculation of N flux from  $^{14}\text{C}$  assumes that C and N entered the plant (or plant and fungus) together without extracellular separation of C and N.  $^{13}\text{C}$  data did not account for respiratory losses and are not shown.

**Figure 5**  $^{15}\text{N}$  distribution within *D. antarctica* roots with and without DSE colonisation after 5 min incubation in either D or L enantiomers of  $^{15}\text{N}$  trialanine. a. Optical image of partial DSE-inoculated root cross-section showing typical cell zonation, specifically the cortex (white inset square) from where nanoSIMS images (c) are taken; scale bar 100  $\mu\text{m}$ . b. TEM of intercellular space between root cortical cells of a DSE-inoculated root showing the presence of abundant hyphae (white arrows); scale bar 2  $\mu\text{m}$ . c. The  $^{15}\text{N}$  atom percent images (nanoSIMS) of typical cortical cells in roots with or without DSE and incubated with either D or L forms of  $^{15}\text{N}$  trialanine. Highest  $^{15}\text{N}$  enrichment was observed in DSE colonised roots supplied with L-trialanine. White arrows indicate intercellular hyphae where they can be clearly identified. Cells of DSE colonised roots supplied with D-trialanine also showed enrichment, but hyphae could not be located with confidence. Roots without DSE showed negligible  $^{15}\text{N}$  enrichment; scale bar 10  $\mu\text{m}$ .




**Figure 1**




**Figure 2**






**Figure 3**



**Figure 4**

514



515

516 **Figure 5**

517

518

519

520

521



522

523

524 **Figure S1** Moss carpet dominated by *Sanionia uncinata* on Signy Island.

525

526

527

528



529



530

531

532 **Figure S2** Moss banks on Signy Island, showing living *Chorisodontium aciphyllum* with  
533 accumulated moss peat underneath.

534

535

536



537

538


539

540

541

542

**Figure S3** *Deschampsia antarctica* with shallow roots penetrating into accumulated organic matter under *Sanionia uncinata*. Pencil gives scale (c. 6 mm diameter).




543

544 **Figure S4** Location of Signy Island (inset) in maritime Antarctica. Plants were sampled from (1)

545 North Point, (2) Deschampsia Point, (3) Moss Braes, (4) Starfish Cove, (5) Foca Cove, (6)

546 Berntsen Point, (7) Factory Cove, (8) Factory Bluffs, (9) Polynesia Point and (10) Gourlay Point.

547



548


549

550 **Figure S5** Apparent rates of uptake of C from D-and L-enantiomers of alanine and short peptides  
 551 thereof into shoots of *D. antarctica* and *C. quitensis* following injection of <sup>15</sup>N- and <sup>13</sup>C- labelled  
 552 substrates into soil. Values are mean ± SEM; *n*=3 or 4. Caution should be exercised in  
 553 interpretation as differences in partitioning and losses of <sup>13</sup>C in respiration are not accounted for.


554

555






**Figure S6** Ratio of recovery of isotopic labels in roots to recovery in shoots of *D. antarctica*. a.  $^{14}\text{C}$ . b.  $^{13}\text{C}$ . c.  $^{15}\text{N}$ . Data are mean  $\pm$  SEM;  $n=3$  for  $^{14}\text{C}$ ;  $n=4$  for  $^{13}\text{C}$  and  $^{15}\text{N}$ .



**Figure S7** Loss in respiration of C supplied to roots of *D. antarctica* as D- and L-alanine and their short peptides. Closed and open circles are plants colonised with DSE and uncolonulated controls, respectively. Data are mean  $\pm$  SEM;  $n=3$ .



565

566 **Figure S8**  $^{15}\text{N}$  enrichment image of entire root cross-section of *D. antarctica* inoculated with  
567 DSE and incubated for 5 min in  $^{15}\text{N}$  L-trialanine. Similar montaged images were generated for all  
568 four treatments (+/-DSE, D or L  $^{15}\text{N}$ -trialanine) and an unlabelled control. For the example  
569 displayed here (A),  $^{15}\text{N}$  enrichment is highest in the intercellular spaces of the cortical zone and  
570 also in portions of microscleotia. B)  $^{12}\text{C}^{14}\text{N}^-$  ion (proxy for  $^{14}\text{N}$ ) intensity image of the same area  
571 is included as a reference to sample ultrastructure.

572 **Table S1** Michaelis-Menten constants for uptake of various forms of N by roots of *D. antarctica*  
573 without or with DSE colonisation

	Km ( $\mu\text{mol l}^{-1}$ )		Vmax ( $\mu\text{mol g}^{-1} \text{DW root h}^{-1}$ )		Difference between DSE colonised and control plants
	-DSE	+DSE	-DSE	+DSE	
NO <sub>3</sub> <sup>-</sup>	3488	3308	42.0	36.8	<i>P</i> =0.90
NH <sub>4</sub> <sup>+</sup>	5743	5191	35.9	42.2	<i>P</i> =0.007
L-alanine	323.1	840.6	12.5	42.7	<i>P</i> =0.07
D-alanine	657.1	782.3	15.2	20.6	<i>P</i> =0.04
L-dialanine	222.2	410.9	11.1	45.3	<i>P</i> =0.02
D-dialanine	1261	663.4	27.0	23.2	<i>P</i> =0.08

574

575

576